

(12) DEMANDE INTERNATIONALE PUBLIÉE EN VERTU DU TRAITÉ DE COOPÉRATION  
EN MATIÈRE DE BREVETS (PCT)

(19) Organisation Mondiale de la Propriété  
Intellectuelle  
Bureau international



(43) Date de la publication internationale  
12 avril 2001 (12.04.2001)

PCT

(10) Numéro de publication internationale  
WO 01/25227 A2

(51) Classification internationale des brevets:  
C07D 401/00

(71) Déposant: AVENTIS PHARMA S.A. [FR/FR]; 20 av-  
enue Raymond Aron, F-92160 Antony (FR).

(21) Numéro de la demande internationale:  
PCT/FR00/02541

(72) Inventeurs: MALLERON, Jean-Luc; 2 allée Renoir,  
F-91460 Marcoussis (FR). TABART, Michel; 3 rue  
Paul Langevin, F-91290 La Norville (FR). CARRY,  
Jean-Christophe; 5 rue de Savoie, F-94100 Saint Maur  
(FR). EVERS, Michel; 8 rue Robert Schumann, F-94510  
La Queue En Brie (FR). EL AHMAD, Yousef; 11 avenue  
de Verdun, F-94000 Creteil (FR). MIGNANI, Serge; 14  
avenue de Robinson, F-92290 Chatenay Malabry (FR).  
VIVIANI, Fabrice; 46 rue Jules Fossier, F-95380 Louvres  
(FR).

(22) Date de dépôt international:  
14 septembre 2000 (14.09.2000)

(25) Langue de dépôt: français


(26) Langue de publication: français

(30) Données relatives à la priorité:  
99/11679 17 septembre 1999 (17.09.1999) FR

[Suite sur la page suivante]

(54) Title: PIPERIDINE QUINOLYL PROPYL DERIVATIVES, PREPARATION METHOD AND COMPOSITIONS CONTAIN-  
ING SAME

(54) Titre: DERIVES DE LA QUINOLYL PROPYL PIPERIDINE, LEUR PREPARATION ET LES COMPOSITIONS QUI LES  
CONTIENNENT



(57) Abstract: The invention concerns piperidine quinolyl propyl derivatives of general formula (I) wherein : R<sub>1</sub> is H or halogen, or OH, R'<sub>1</sub> is H, or may represent halogen when R<sub>1</sub> is also halogen, and R\* is H, or R<sub>1</sub> and R\* together form a bond and R'<sub>1</sub> is H, R<sub>2</sub> is a carboxy, carboxymethyl or carboxy-2-ethyl radical, and R<sub>3</sub> is C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 1 to 3 substituents selected among OH, halogen, =O, COOH, alkyloxycarbonyl, alkyloxy, alkylthio or among a phenyl, phenylthio or phenylalkylthio radical which can themselves bear 1 to 4 substituents [selected among halogen, OH, alkyl, alkyloxy, trifluoromethyl, trifluoromethoxy, COOH, alkyloxycarbonyl, CN, acetamido or NH<sub>2</sub>] or among cycloalkyl, cycloalkylthio (3 to 7 members), or among aromatic heterocyclyl or heterocyclylthio (5 to 6 members) comprising 1 to 4 heteroatoms selected among N, O or S and optionally substituted themselves [with halogen, OH, alkyl, alkyloxy, CF<sub>3</sub>, OCF<sub>3</sub>, =O, COOH, alkyloxycarbonyl, CN or NH<sub>2</sub>], R<sub>3</sub> is propargyl substituted with phenyl which can itself bear 1 to 3 substituents [selected among halogen, OH, alkyl, alkyloxy, CF<sub>3</sub>, OCF<sub>3</sub>, COOH, alkyloxycarbonyl, CN or NH<sub>2</sub>], or substituted with a cycloalkyl radical comprising 3 to 7 members or substituted with aromatic heterocycle (5 to 6 members) comprising 1 to 4 heteroatoms selected among N, O or S and optionally substituted itself [with halogen, OH, alkyl, alkyloxy, CF<sub>3</sub>, OCF<sub>3</sub>, =O, COOH, alkyloxycarbonyl, CN or NH<sub>2</sub>], or R<sub>3</sub> represents cinnamyl or 4-phenylbuten-3-yl, or R<sub>2</sub> is -CH<sub>2</sub>OH, alkyloxy-carbonyl, alkyloxycarbonylmethyl or alkyloxycarbonyl-2-ethyl and R<sub>3</sub> is C<sub>1</sub>-C<sub>6</sub> alkyl substituted with phenylthio which can itself bear 1 to 3 substituents [selected among halogen, OH, alkyl, alkyloxy, CF<sub>3</sub>, OCF<sub>3</sub>, COOH, alkyloxycarbonyl, CN or NH<sub>2</sub>], with cycloalkylthio comprising 3 to 7 members, or with aromatic heterocyclylthio (5 to 6 members) comprising 1 to 4 heteroatoms selected among N, S or O and optionally substituted itself [with halogen, OH, alkyl, alkyloxy, CF<sub>3</sub>, OCF<sub>3</sub>, =O, COOH, alkyloxycarbonyl, CN or NH<sub>2</sub>] or R<sub>3</sub> is propargyl substituted with phenyl, which can itself bear 1 to 3 substituents [selected among halogen OH, alkyl, alkyloxy, CF<sub>3</sub>, OCF<sub>3</sub>, COOH, alkyloxycarbonyl, CN or NH<sub>2</sub>], or substituted with cycloalkyl comprising 3 to 7 members or substituted with an aromatic heterocyclyl with 5 to 6 members comprising 1 to 4 heteroatoms selected among N, O or S and optionally substituted itself [with halogen, OH, alkyl, alkyloxy, CF<sub>3</sub>, OCF<sub>3</sub>, =O, COOH, alkyloxycarbonyl, CN or NH<sub>2</sub>], and R<sub>4</sub> is C<sub>1</sub>-C<sub>6</sub> alkyl, (C<sub>3</sub>-C<sub>7</sub>) alkenyl-CH<sub>2</sub>- or alkynyl-CH<sub>2</sub>.

[Suite sur la page suivante]